

# Hyperbolicity of GR in null foliations

Thanasis Giannakopoulos

Instituto Superior Técnico, Lisbon


GWverse COST action meeting, Lisbon, September 1, 2021



---

PhysRevD.102.064035 with D. Hilditch & M. Zilhão  
work in progress with N. Bishop, D. Hilditch, D. Pollney & M. Zilhão

# Highly accurate gravitational waveform modelling



Cauchy-Characteristic extraction

# Hyperbolicity

$$\mathcal{A}^t(\mathbf{u}, x^\mu) \partial_t \mathbf{u} + \mathcal{A}^p(\mathbf{u}, x^\mu) \partial_p \mathbf{u} + \mathcal{S}(\mathbf{u}, x^\mu) = 0,$$

where  $\mathbf{u} = (u_1, u_2, \dots, u_q)^T$ , is the state vector of the system and

$$\mathcal{A}^\mu = \begin{pmatrix} a_{11}^\mu & \cdots & a_{1q}^\mu \\ \vdots & \ddots & \vdots \\ a_{q1}^\mu & \cdots & a_{qq}^\mu \end{pmatrix}$$

denotes the principal part matrices, with  $\det(\mathcal{A}^t) \neq 0$ . Construct the

$$\mathbf{P}^s = (\mathcal{A}^t)^{-1} \mathcal{A}^p s_p,$$

where  $s^i$  is an arbitrary unit spatial vector.


## Well-posedness

The PDE problem has a unique solution that depends continuously on the given data in a suitable norm.

- Strongly hyperbolic (SH) → **well-posed** IVP in the  $L^2$  norm
- Weakly hyperbolic (WH) → **ill-posed** IVP in the  $L^2$  norm

A numerical solution **can converge** to the continuum **only** for well-posed PDE problems.

## Bondi-like gauges




- vector basis:  $\partial_u, \partial_r, \partial_\theta, \partial_\phi$
- $\partial_r$  is null &  $\perp$  to  $\partial_\theta$  and  $\partial_\phi$

$$g^{\mu\nu} = \begin{pmatrix} 0 & g^{ur} & 0 & 0 \\ g^{ur} & g^{rr} & g^{r\theta} & g^{r\phi} \\ 0 & g^{r\theta} & g^{\theta\theta} & g^{\theta\phi} \\ 0 & g^{r\phi} & g^{\theta\phi} & g^{\phi\phi} \end{pmatrix}$$

Vacuum Einstein's equations:

$$\text{Evolution system: } R_{rr} = R_{r\theta} = R_{r\phi} = R_{\theta\theta} = R_{\theta\phi} = R_{\phi\phi} = 0$$

# Weak hyperbolicity of Bondi-like gauges



The principal symbol<sup>1,2</sup>:

$$\mathbf{P}^s = \begin{pmatrix} \boxed{\mathbf{P}_G} & \mathbf{P}_{GC} & \mathbf{P}_{GP} \\ 0 & \mathbf{P}_C & 0 \\ 0 & 0 & \mathbf{P}_P \end{pmatrix}$$

New result<sup>2</sup>:  $\mathbf{P}_G$  is non-diagonalizable along  $\theta$  if  $\partial_r$  is  $\perp$  to  $\partial_\theta$ .

GR in Bondi-like gauges  $\rightarrow$  WH 2nd order PDE system<sup>3</sup>

---

<sup>1</sup>Hilditch & Richter 2016


<sup>2</sup>WIP with Bishop, Hilditch, Pollney & Zilhão

<sup>3</sup>see Ripley 2021 for a symmetric hyperbolic formulation with higher derivatives


# Convergence tests in the $L^2$ norm

- Monitor the numerical error with increasing resolution
- Convergence factor:  $Q = 4$  for these tests by construction


SH smooth




WH smooth



SH noisy



WH noisy



## Summary

- GR in Bondi-like gauges  $\rightarrow$  weakly hyperbolic 2nd order PDE system
- Ill-posed characteristic initial boundary value problem in the  $L^2$  norm (other norms?)
- Weak hyperbolicity in numerics  $\rightarrow$  high frequency given data

### TODO:

- Characteristic GR formulations  $\rightarrow$  strongly hyperbolic 2nd order PDE system



## Summary

- GR in Bondi-like gauges  $\rightarrow$  weakly hyperbolic 2nd order PDE system
- Ill-posed characteristic initial boundary value problem in the  $L^2$  norm (other norms?)
- Weak hyperbolicity in numerics  $\rightarrow$  high frequency given data

### TODO:

- Characteristic GR formulations  $\rightarrow$  strongly hyperbolic 2nd order PDE system

Thank you!